

The Wouthuysen-Field effect in a clumpy intergalactic medium

Jonathan Higgins, Avery Meiksin

SUPA, Institute for Astronomy, University of Edinburgh, Blackford Hill, Edinburgh EH9 3HJ, UK*

7 November 2008

ABSTRACT

We show that the high optical depth of the intergalactic medium to Ly α photons before the Epoch of Reionization results in a negligible magnitude for the Wouthuysen-Field effect produced by a radiation source on its distant surroundings, unless (1) the scattering medium has sufficient time for the impinging resonance line photons to establish a steady-state frequency distribution or (2) the scattering gas is undergoing internal expansion or has a peculiar motion of tens to hundreds of kms^{-1} away from the source. Because of intergalactic attenuation, discrete structures will receive only radiation from a source displaced from the resonance line frequency by typically hundreds to thousands of Doppler widths. The incident radiation must diffuse across the resonance line to produce a substantial scattering rate. We present steady-state solutions in the radiative diffusion approximation for the radiation field trapped in a clump of gas and show that this may result in an *enhancement* of the strength of the Wouthuysen-Field effect by as much as a factor of 10^6 over the free-streaming (single-scattering) limit. Solutions to the time-dependent diffusion equation, however, show that the timescales required to establish a steady state will generally exceed the lifetime of the sources, resulting in a substantially reduced scattering rate. In the presence of internal expansion, a steady state may be established as photons are redshifted across the resonance line and into the red wing, and significant enhancement in the scattering rate over the free-streaming limit may again be produced. Alternatively, a substantial scattering rate may arise in systems with a peculiar motion away from the source that redshifts the received radiation into the resonance line centre. As a consequence, at epochs $z \lesssim 30$, when collisional decoupling of the hyperfine structure of hydrogen from the Cosmic Microwave Background is small except in dense regions, and prior to the establishment of any large-scale diffuse radiation field of resonance line photons, the 21cm signature from the Intergalactic Medium produced by the Wouthuysen-Field effect will in general trace the peculiar velocity field of the gas in addition to its density structure.

Key words: atomic processes – cosmology: theory – line: formation – radiative transfer – radio lines: general – scattering

1 INTRODUCTION

Following the Recombination Era at $z \simeq 1100$, the baryons produced in the Big Bang were largely neutral. By $z \gtrsim 6$, the spectra of high redshift Quasi-Stellar Objects (QSOs) show that the hydrogen had become ionized (Becker & et al. 2001). The history of structure formation in the wide redshift expanse between these epochs, when there were few if any sources of radiation, is largely unknown. In principle, 21cm emission from intergalactic hydrogen could reveal the evolution of structure in the baryons during these cos-

mic “Dark Ages.” Because the baryons are cold, they would closely trace the evolution of the dark matter, so that 21cm imaging could trace the growth of structure following the Recombination Era (Hogan & Rees 1979; Scott & Rees 1990). Because of strong coupling between the hyperfine structure of the hydrogen and the Cosmic Microwave Background (CMB), however, a mechanism that decouples the spin temperature of the hydrogen hyperfine structure from the CMB temperature must be active, otherwise the hydrogen is rendered invisible: it absorbs and re-emits the CMB radiation at the same rate, leaving it indistinguishable from the CMB. At redshifts $z > 30$, collisions between hydrogen and electrons and other hydrogen atoms are adequate to begin de-

* Scottish Universities Physics Alliance

coupling the spin temperature from the CMB (Scott & Rees 1990; Madau et al. 1997). Except in dense regions, however, at later times collisional decoupling is inadequate.

Once the first sources of radiation begin to turn on, the scattering of Ly α photons by the neutral hydrogen offers an alternative means of decoupling, through the Wouthuysen-Field effect (WFE) (Wouthuysen 1952; Field 1958). As sources turn on in sufficient number to begin reionizing the Intergalactic Medium (IGM), they will produce a combined intensity of Ly α photons sufficient for coupling the spin temperature to the light temperature of the Ly α photons rather than the CMB, rendering the intergalactic hydrogen visible against the CMB in either emission or absorption (Madau et al. 1997).

The discovery of the End of the Dark Ages (EDA) and the onset of the Epoch of Reionization (EoR) has become one of the paramount goals of a new generation of radio telescopes, such as the LOw Frequency Array (LOFAR)¹, the Murchison Widefield Array (MWA)², the Primeval Structure Telescope/21 Centimeter Array (PaST/21CMA)³, the Precision Array to Probe EoR (PAPER)⁴, and a possible Square Kilometre Array (SKA)⁵. Reviews of this rapidly growing area are provided by Loeb & Barkana (2001), Fan et al. (2006) and Furlanetto et al. (2006).

Madau et al. (1997) estimated the Ly α collision rate P_α that drives the WFE as the integrated intensity from cosmologically distributed sources, assuming that photons blueward of Ly α emitted from a source will contribute their full amount to P_α once they redshift into the local Ly α resonance. In fact, this provides a lower limit (assuming photons are not destroyed by, eg, dust absorption). The multiple scattering of resonance line photons will in general enhance the radiation field. For pure Doppler scattering, Field (1959a) argued the rate is enhanced by the number of scatters a photon undergoes before being randomly scattered sufficiently redward of line centre to escape. The estimate neglected scattering in the Lorentz wings, however. It also assumed that the IGM takes part in the homogeneous and isotropic expansion of the Universe. In fact the IGM is clumpy, with structures breaking away from the cosmological expansion. Efforts are underway to estimate the radiation field and scattering rate in a cosmological context through Monte Carlo simulations of the scattering of resonance line photons in an inhomogeneous medium (Chuzhoy & Zheng 2007; Pierleoni et al. 2007; Semelin et al. 2007). In this paper, we use analytical means to explore some of the consequences of intergalactic structure formation for the WFE as a means of generating an intergalactic 21cm signature.

In the next section, we discuss the optical depth of the IGM to resonance line photons and the implications for the scattering rate. Approximate steady-state solutions to the radiative transfer equation are derived in § 3, and time-dependent solutions in § 4. A discussion of the solutions and applications are provided in § 5, along with a summary of our conclusions.




Figure 1. (Upper panel) The optical depth of photons emitted by a source at redshift z_S and received at the frequency ν near the resonance line frequency ν_0 of Ly α at redshift z . The frequency is expressed as $x = (\nu - \nu_0)/\Delta\nu_D$. The curves are for $T = 10$ K (solid), 100 K (dashed) and 1000 K (dot-dashed). Also shown is the approximation $\tau_\nu \simeq x_1/x$ for the $T = 10$ K model (dotted line, nearly coinciding with the solid line). (Lower panel) The value of x at which $\tau_\nu = 1$ for $z_S - 1 < z < z_S$. The curves are labelled as in the upper panel.

2 THE WFE IN THE INTERGALACTIC MEDIUM

The optical depth through a homogeneous and isotropic expanding IGM of a photon emitted by a source at redshift z_S and received at redshift z at frequency $\nu > \nu_0$, where ν_0 is the resonance line frequency, is (Field 1959a; Gunn & Peterson 1965)

$$\tau_\nu = \sigma \int_z^{z_S} dz' \frac{dl_p}{dz'} n_l(z') \varphi_V \left(a, \nu \frac{1+z'}{1+z} \right), \quad (1)$$

where $n_l(z')$ is the number density of scattering atoms in the lower level at epoch z' , $\sigma = \pi e^2 f_{lu}/(m_e c) \simeq 0.0110 \text{ cm}^2 \text{ Hz}$ is the total resonance line cross section, where $f_{lu} \simeq 0.4162$ is the upward oscillator strength for hydrogen Ly α , $\varphi_V(a, \nu)$ is the Voigt line profile normalized to $\int d\nu \varphi_V(a, \nu) = 1$, $a \simeq 0.0472 T^{-1/2}$ is the ratio of the decay rate to the Doppler width $\Delta\nu_D = \nu_0 b/c$, where $b = (2k_B T/m_H)^{1/2}$ is the Doppler parameter for hydrogen gas at temperature T and c is the speed of light, and l_p is the proper path length. In the Lorentz wing, expressed as a function of $x = (\nu - \nu_0)/\Delta\nu_D$, the dimensionless Voigt profile $\phi_V(a, x) = (\Delta\nu_D) \varphi_V(a, \nu)$ is well-approximated as $\phi_V(a, x) \simeq a/(\pi x^2)$. The differential proper line element evolves according to $dl_p/dz \simeq (c/H_0) \Omega_m^{-1/2} (1+z)^{-5/2}$ in a flat universe at redshifts for which $\Omega_m (1+z)^3$ dominates the contribution from the vacuum energy, where Ω_m is the ratio of the total mass density to the critical Einstein-deSitter density, and where $H_0 = 100h \text{ km s}^{-1} \text{ Mpc}^{-1}$ is the Hubble constant today.

¹ www.lofar.org

² www.haystack.mit.edu/ast/arrays/mwa

³ web.phys.cmu.edu/~past

⁴ astro.berkeley.edu/~dbacker/eor

⁵ www.skatelescope.org

Photons emitted by the source at a frequency $\nu_e > \nu_0(1+z_S)/(1+z)$ will scatter in the blue Lorentz wing (except for those for which the inequality is nearly an equality, in which case they will scatter through the Doppler core). In the blue wing the optical depth due to Ly α scattering by neutral hydrogen is

$$\tau_\nu \simeq 1.75 \times 10^5 \frac{cn_i(0)}{H_0 \Omega_m^{1/2}} \frac{(1+z)^{3/2}}{\nu_0^2 y^{3/2}} \left\{ \left[\frac{y^{1/2}}{y-1} - \frac{u^{1/2}}{u-1} \right] + \frac{1}{2} \log \left[\frac{(1-u^{1/2})(1+y^{1/2})}{(1-y^{1/2})(1+u^{1/2})} \right] \right\}, \quad (2)$$

where $y = \nu/\nu_0$ and $u = y(1+z_S)/(1+z)$, and all quantities are assumed to be in cgs units. For $0 < xb/c < (z_S - z)/(1+z) < 1$, the expression simplifies to

$$\tau_\nu \approx \frac{x_1}{x}, \quad (3)$$

where

$$x_1 = \frac{a}{\pi} \frac{\sigma c}{\nu_0} \frac{n_i(0)}{H_0 \Omega_m^{1/2}} (1+z)^{3/2} \approx 214 h^{-1} (1+z)^{3/2} T_{\text{IGM}}^{-1/2} \quad (4)$$

for a mean line-of-sight IGM temperature T_{IGM} for a universe with $\Omega_m = 0.3$ and baryon density $\Omega_b h^2 = 0.022$ (O'Meara et al. 2006; Spergel & et al. 2007). As an illustration, $\tau_\nu(x)$ for a source at $z_S = 8$ is shown at $z = 7$ in Fig. 1 (upper panel). The optical depth is extremely high until ν is well displaced from the line centre. The values of x_1 , at which $\tau_\nu = 1$, are shown in the lower panel of Fig. 1 for a range of redshifts.

Photons emitted by the source with frequencies between $\nu_0 < \nu_e < \nu_0(1+z_S)/(1+z)$ will pass through the resonance line frequency en route to the gas at z . Upon passing through the resonance line, the photons will essentially be completely scattered out of the line of sight. As a consequence, essentially no photons will be received in the frequency range $\nu_0(1+z)/(1+z_S) < \nu < \nu_0$ (the Gunn-Peterson effect). A general expression for the optical depth through the Doppler core is provided in Appendix A.

Photons emitted sufficiently redward of ν_0 to avoid the Doppler core may still be scattered in the Lorentz red wing of the surrounding hydrogen. This will produce a halo of scattered Ly α photons around a central source like a QSO (Loeb & Rybicki 1999). For frequencies $x < 0$ near line centre, Loeb & Rybicki (1999) show that the photons will build up an energy density as they diffuse through the surrounding IGM of the form $u_\nu \sim (-x)^{-9/2} e^{-[r/r_c(x)]^2}$, where $r_c(x) = (-x)^{1/2} (2/3) [(b/H) \lambda_{\text{mfp}}(x)]^{1/2}$ scales like the harmonic mean between the distance over which Hubble expansion produces a velocity difference matching the Doppler parameter b and the scattering mean free path of a resonance line photon in the Lorentz wing

$$\lambda_{\text{mfp}}(x) \simeq \frac{\pi \Delta \nu_D}{n_i \sigma a} x^2, \quad (5)$$

where n_i is the number density of atoms in the lower level. The energy density is exponentially suppressed for $x \rightarrow 0^-$ at a fixed position, resulting in a negligible Ly α scattering rate except very near the QSO. Numerically,

$$r_c(x) \simeq 0.6 \text{ pc} (-x)^{3/2} h^{-1/2} T_{\text{IGM}}^{3/4} \Omega_m^{-1/4} \left(\frac{1+z}{9} \right)^{-9/4}, \quad (6)$$

so that a substantial Ly α scattering rate is confined to a small region around the source. The region is so small in fact, that the assumption that the surrounding IGM takes part in the Hubble flow will not be valid. More likely the gas will have been ionized and violently disturbed by the QSO itself.

Finally, photons emitted sufficiently redward of ν_0 at the source to escape scattering in the Lorentz wing will pass freely through the IGM, where they will be received at redshift z at frequencies $x < -x_0$, where

$$x_0 = \frac{c}{b_{\text{IGM}}} \frac{z_S - z}{1 + z_S} \simeq 2.3 \times 10^6 T_{\text{IGM}}^{-1/2} \frac{z_S - z}{1 + z_S}. \quad (7)$$

The vast majority of photons from the source emitted blueward of, but near, the resonance line frequency never arrive near the line centre, having been scattered out of the path along the way. The resonance line radiation impinging on the absorption site⁶ from the source scatters in the Ly α resonance line at the rate per neutral atom in the lower state

$$P_l = \sigma c \int_0^\infty d\nu \varphi_V(a, \nu) \frac{u_\nu}{h_P \nu}, \quad (8)$$

where h_P is Planck's constant and u_ν is the local specific energy density of photons received at z from the source at z_S with specific luminosity L_ν

$$u_\nu = \frac{1}{c} \frac{L_\nu}{4\pi r_L^2} \exp(-\tau_\nu) \quad (9)$$

where r_L is the luminosity distance between the source and the gas at redshift z . The scattering rate assuming none of the photons have been scattered out of the line of sight en route to the absorption site (the free-streaming limit) is

$$P_l(\tau_\nu = 0) \simeq \sigma \frac{L_\nu / h_P \nu}{4\pi r_L^2}. \quad (10)$$

The ratio of the scattering rate for $\tau_\nu \simeq x_1/x$ to the free-streaming value is then given by

$$\frac{P_l(\tau_\nu)}{P_l(\tau_\nu = 0)} \simeq \frac{\int_0^\infty dx a \pi^{-1} x^{-2} \exp(-x_1/x)}{\int_{-\infty}^\infty dx \phi_V(a, x)} = \frac{a}{\pi x_1} \simeq 7.0 \times 10^{-5} h (1+z)^{-3/2} \left(\frac{T_{\text{IGM}}}{T_a} \right)^{1/2}, \quad (11)$$

where T_a is the temperature of the absorbing gas. The low value poses a fundamental problem to the effectiveness of the WFE as a means of decoupling the spin state of the gas from statistical equilibrium with the CMB. *Only if photons have sufficient time to diffuse across the line centre will the scattering of Ly α photons be an effective means of decoupling the spin temperature from the CMB in comoving objects.*

3 THE WFE IN DISCRETE OBJECTS

3.1 Steady-state solutions without atomic recoil

The assumption of a homogeneous and isotropic universe, while a fair approximation over large scales, breaks

⁶ Strictly speaking, the photons are scattered. The region at which the scattering rate is computed is referred to for simplicity as the absorption site or the absorber.

down on small scales: the IGM is clumpy at the redshifts from which the 21cm signal is expected (Tozzi et al. 2000; Gnedin & Shaver 2004). For a sufficiently short mean free path, photons will be trapped within a discrete system and diffuse in frequency and space as they scatter within the system. In this section, we consider the energy distribution the radiation will reach given adequate time to relax to a steady state.

For a Ly α photon to become trapped within an absorber of lengthscale L_a , the mean free path must minimally satisfy $\lambda_{\text{mfp}} < L_a$, for which $\tau_\nu = L_a/\lambda_{\text{mfp}} > 1$. As the optical depth in the wing may be expressed as $\tau_\nu = \pi^{-1/2}(a\tau_0)x^{-2}$, where $\tau_0 = n_i L_a \pi^{-1/2} \sigma(\Delta\nu_D)^{-1}$ is the optical depth at the line centre, this corresponds to photons escaping with $x > \pi^{-1/4}(a\tau_0)^{1/2}$ (Osterbrock 1962). Radiative transfer solutions for sources embedded within a slab suggest this may be somewhat too restrictive for systems with very high line centre optical depths, in which case the photons escape through spatial diffusion rather than frequency diffusion. Steady-state solutions in the diffusion approximation for scattering in the Lorentz wings in very high optical depth systems show that photons escape the slabs with frequencies typically on the order of $(a\tau_0)^{1/3}$ (Adams 1972; Harrington 1973; Neufeld 1990). An escape frequency of $x_{\text{esc}} = x_*(a\tau_0)^{1/3}$ corresponds to a mean free path smaller than the thickness of the slab by the factor $f = x_*^2 \pi^{1/2} (a\tau_0)^{-1/3} \simeq 0.013 x_*^2 T_a^{1/3} N_{19}^{-1/3}$, where T_a is the temperature of the absorber and N_{19} is the column density in units of 10^{19} cm^{-2} . (This corresponds to a region 24 kpc across for the mean intergalactic hydrogen density at $z = 8$, and smaller for overdense systems.) As f depends only weakly on T_a and N_{HI} , while x_* depends on $a\tau_0$ and the radiation source parameters of any particular problem, we express the escape frequency simply as

$$x_{\text{esc}} \simeq 395(10f)^{1/2} N_{19}^{1/2} T_a^{-1/2}, \quad (12)$$

noting that f will in general vary with the depth within the absorber, and must be solved for according to each particular configuration, but will be of order 0.01 – 1 for applications we consider. Comparison of Eq. (12) with Eq. (4) for x_1 shows that photons with $\tau_\nu < 1$ in the IGM will generally not be trapped within a structure comoving with the expansion of the Universe unless the structure has a neutral hydrogen column density $N_{19} > N_{19,\text{crit}}$, where

$$N_{19,\text{crit}} \simeq 0.29h^{-2}(10f)^{-1}(1+z)^3 \left(\frac{T_a}{T_{\text{IGM}}} \right). \quad (13)$$

If $x_1 < x_{\text{esc}}$, the absorber will behave as a photon bucket, allowing the energy density of the radiation field at frequencies $x < x_{\text{esc}}$ to build up with time as radiation from the source becomes trapped. As the photons scatter within the absorber, they will diffuse in frequency and space. Because photons with $|x| > x_{\text{esc}}$ will escape, the energy density of the radiation trapped within the absorber will reach a steady state when the rate of incoming photons balances the rate of escape. In principle, the energy density that develops could produce a substantial scattering rate, even exceeding the rate estimated assuming no photon losses from the source due to scattering by the intervening IGM.

The depth to which photons at frequency x_1 penetrate an absorber depends on the optical depth of the absorber. If it is optically thin at x_1 , the photons will stream through.

If optically thick, the photons will scatter near the surface of the absorber. Photons blueward of x_1 , however, will penetrate more deeply, as the cross section diminishes like $1/x^2$. In general, photons will be injected at varying layers within the absorber, with bluer photons injected at increasing depths.

We would like to estimate the evolution of the radiation field within an absorber. No solution to this problem exists in the literature. The full radiative transfer problem through the slab is an involved one, as the radiation field will vary with both the optical depth through the slab and with frequency. It is particularly important in our application to obtain the solution across the Doppler core to estimate the Ly α scattering rate. Instead of solving the full problem, we seek approximate solutions in which we treat the scattered radiation field as locally isotropic, neglecting the spatial diffusion of the radiation through the absorber. We show in Appendix B that the resulting radiative transfer equation in the diffusion approximation is formally identical to the equation of Harrington (1973) for the problem of a slab with a uniform distribution of spectrally flat sources. The solution we obtain neglecting the spatial diffusion of the radiation agrees well with Harrington’s solution which includes both spatial and frequency diffusion. A detailed quantitative description will require a more exact solution to the problem, but our approximate approach allows us to explore qualitatively several effects of interest. We would like to estimate the energy density that may be achieved as a function of x_{esc} . We also wish to examine the effect of internal motions on the radiation energy density across the line centre and the possible role of atomic recoil. Lastly, we would like to estimate the time it takes the radiation field to establish a steady state. All of these effects extend well beyond those explored by existing slab solutions.

We estimate the radiation density that builds up within an absorber by assuming a steady state between the rate of photons injected into the absorber and the rate of diffusion of the photons across the escape frequency, imposing the boundary condition $u(x) = 0$ for $|x| > x_{\text{esc}}$, where $u(x) = \Delta\nu_D u_\nu$. We assume isotropic scattering within a homogeneous absorber, and use the diffusion approximation to describe the resonance line scattering of the photons. In equilibrium,

$$\frac{1}{2} \frac{d}{dx} \left[D(a, x) \frac{dn(x)}{dx} \right] = -\tilde{S}(x), \quad (14)$$

where $n(x) = \Delta\nu_D n_\nu$, $n_\nu = u_\nu/h\nu$, $\tilde{S}(x) = (\Delta\nu_D)^2 S(\nu)/(n_i c \sigma)$, where $S(\nu)$ is a source function, and $D(a, x)$ is the diffusion coefficient. The second order moment of the frequency redistribution function gives for the diffusion coefficient $D(a, x) = \phi_V(a, x) + (1/3)d^2\phi_V(a, x)/dx^2$ (Rybicki & dell’Antonio 1994). The derivation of the diffusion equation, however, does not conserve photon number to the order of the approximation. As a consequence, the diffusion coefficient is left ambiguous. Rybicki & dell’Antonio (1994) advocate adopting $D(a, x) = \phi_V(a, x)$ for its simplicity, and we shall do so here.

The source $S(\nu)$ represents the rate at which photons received from the source are incident on the absorber at $x < x_{\text{esc}}$, where they are trapped and diffuse in frequency. It is given by the scattering emissivity




Figure 2. The steady-state distribution of Ly α resonance line photons for a source function $\tilde{S}(x) = \delta_D(x - x_{\text{inj}})$ with $x_{\text{inj}} = 100$, allowing photons with $|x| > x_{\text{esc}} = 200$ to escape, as given by the diffusion approximation, Eq. (14), with $D(a, x) = \phi_V(a, x)$. An absorber temperature $T_a = 100$ K is assumed.

$$S(\nu) = n_l c \sigma \int_0^\infty d\nu' R(\nu', \nu) n_{\nu'}, \quad (15)$$

where $R(\nu', \nu)$ is the photon redistribution function for resonance line scattering (Mihalas 1978). Since scattering in the wings is coherent, $R(\nu', \nu) \simeq \varphi_V(a, \nu') \delta_D(\nu' - \nu)$, where δ_D is the Dirac δ -function. Then

$$\begin{aligned} S(\nu) &\simeq n_l c \sigma \varphi_V(a, \nu) n_\nu \\ &\simeq n_l \sigma a \pi^{-1} (\Delta\nu_D)^{-1} \frac{L_\nu / h_P \nu}{4\pi r_L^2} x^{-2} \exp(-x_1/x). \end{aligned} \quad (16)$$

The dependence on x results in a sharp peak at $x = x_1/2$, so that the source $\tilde{S}(x)$ is well approximated by

$$\begin{aligned} \tilde{S}(x) &\simeq \frac{a}{\pi c} \frac{\Delta\nu_D}{x_1} \frac{L_\nu / h_P \nu}{4\pi r_L^2} \delta_D(x - x_{\text{inj}}) \\ &= A \delta_D(x - x_{\text{inj}}) \end{aligned} \quad (17)$$

where the normalization $A = 2.5 \times 10^{-6} h(1+z)^{-3/2} T_{\text{IGM}}^{1/2} [(L_\nu / h_P \nu) / 4\pi r_L^2]$ (with all quantities expressed in cgs units), matches the source rate of Eq. (16) integrated over all frequencies. Note that the dependence on the absorber temperature cancels. Here, the frequency at which photons are injected into the absorber is represented generally as x_{inj} to allow for any peculiar motion of the absorber, which could shift the peak of the received radiation either redward or blueward, and to allow for scattering within the absorber which will shift the peak of the radiation field penetrating to deeper layers further toward the blue. For a comoving absorber, $x_{\text{inj}} = x_1/2$ near the surface. The solution to Eq. (14) using the wing approximation for $\phi_V(a, x)$ is

$$n(x) = \begin{cases} \frac{\pi}{3a} A \left(1 - \frac{x_{\text{inj}}^3}{x_{\text{esc}}^3}\right) (x_{\text{esc}}^3 + x^3) & ; -x_{\text{esc}} < x < x_{\text{inj}} \\ \frac{\pi}{3a} A \left(1 + \frac{x_{\text{inj}}^3}{x_{\text{esc}}^3}\right) (x_{\text{esc}}^3 - x^3) & ; x_{\text{inj}} < x < x_{\text{esc}}. \end{cases} \quad (18)$$

The solution is shown in Fig. 2. We find that adopting $D(a, x) = \phi_V(a, x) + (1/3)d^2\phi_V(a, x)/dx^2$ produces a nearly identical solution, agreeing to within a fraction of a percent for x_{inj} well out of the core. Numerically integrating the diffusion equation using the full Voigt profile produces a solution that also differs negligibly from Eq. (18).

The ratio of the collision rate P_l , given by using Eq. (18)




Figure 3. The steady-state distribution of Ly α resonance line photons for a source function $\tilde{S}(x) = \delta_D(x - x_{\text{inj}})$ with $x_{\text{inj}} = 100$ in an absorber with temperature $T_a = 100$ K, allowing photons with $|x| > x_{\text{esc}} = 200$ to escape and including the effects of internal motions with a dimensionless velocity gradient $\gamma = 1.7 \times 10^{-7}$ (solid line), representing expansion. Virtually all the injected photons are redshifted, escaping for $x < -x_{\text{esc}}$. The solution for $x_{\text{inj}} = 25$ and $x_{\text{esc}} = 50$ for $\gamma = 1.7 \times 10^{-7}$ is also shown (long-dashed line), showing a weak tail at $x > x_{\text{inj}}$. The corresponding case with contraction, $\gamma = -1.7 \times 10^{-7}$ (short-dashed line) shows a more complex behaviour as contraction curtails the redward diffusion.

in Eq. (8), to the free-streaming collision rate $P_l(0)$ of Eq. (10) is then

$$\begin{aligned} \frac{P_{l,\text{ss}}}{P_l(0)} &\simeq \frac{1}{x_1} \frac{a}{\pi A} \int_{-\infty}^{\infty} dx \phi_V(a, x) n(x) \\ &\simeq \frac{1}{3x_1} \left[\left(1 - \frac{x_{\text{inj}}^3}{x_{\text{esc}}^3}\right) x_{\text{esc}}^3 + \frac{a}{\pi} \left(3x_{\text{inj}}^2 - x_{\text{esc}}^2 - 2\frac{x_{\text{inj}}^3}{x_{\text{esc}}}\right) \right] \\ &\simeq \frac{1}{3x_1} \left(1 - \frac{x_{\text{inj}}^3}{x_{\text{esc}}^3}\right) x_{\text{esc}}^3 \\ &\simeq 9.6 \times 10^4 h \left[\frac{(10f)N_{19}}{(1+z)T_a} \right]^{3/2} T_{\text{IGM}}^{1/2} \left(1 - \frac{x_{\text{inj}}^3}{x_{\text{esc}}^3}\right), \end{aligned} \quad (19)$$

where we have used the approximation Eq. (A1) below for the Voigt profile with $x_{\text{esc}} > x_{\text{inj}} \gg x_m$ assumed, and neglected the terms following the leading. The enhancement may be substantial. For $N_{19} = N_{19,\text{crit}}$, $P_{l,\text{ss}}/P_l(0) \simeq 1.5 \times 10^4 h^{-2} (1+z)^3 T_{\text{IGM}}^{-1} (1 - x_{\text{inj}}^3/x_{\text{esc}}^3)$, so that even for a warm IGM temperature of $T_{\text{IGM}} = 1000$ K a substantial boost in the scattering rate will result.

The solution Eq. (18) assumes velocities internal to the cloud are negligible. To estimate the effect internal motion may have on the frequency distribution of the photons, we approximate the motion as an isotropic expansion (or contraction) within the cloud characterised by a uniform velocity gradient H_a (which in general will not be the same as the Hubble parameter). In the diffusion approximation, Eq. (14) is modified to (Rybicki & dell'Antonio 1994)

$$\frac{1}{2} \frac{d}{dx} \left[2\gamma n(x) + D(a, x) \frac{dn(x)}{dx} \right] = -\tilde{S}(x), \quad (20)$$

where $\gamma = \nu_0 H_a / (\sigma n_l c)$ is (c/b) times the ratio of the scattering time of the photons at line centre to the expansion time H_a^{-1} of the gas. For this solution, the steady state applies to the comoving number density of photons per frequency (Rybicki & dell'Antonio 1994). Overdense struc-




Figure 4. The steady-state distribution of Ly α resonance line photons in the diffusion approximation for a source function $\tilde{S}(x) = \delta_D(x - x_{\text{inj}})$ with $x_{\text{inj}} = 100$, allowing photons with $|x| > x_{\text{esc}} = 200$ to escape and including the effect of atomic recoils. Shown are the solution with $\epsilon = 0.0079$, corresponding to $T_a = 10$ K (solid line), the solution with $\epsilon = 0$ (long-dashed line), and the first order solution $n_0(x) + \epsilon n_1(x)$ for $\epsilon = 0.0079$ (short-dashed line).

tures are typically contracting, but the internal motions along a filament may expand along the filament in regions (Zhang et al. 1998). For an overdensity ten times the cosmic mean at $z = 8$, an internal velocity gradient of 100 km s^{-1} over 100 kpc corresponds to $\gamma \simeq 1.7 \times 10^{-7}$. Contraction by the same magnitude corresponds to $\gamma \simeq -1.7 \times 10^{-7}$.

The solution for the source $\tilde{S}(x) = A\delta_D(x - x_{\text{inj}})$ is

$$n(x) = \begin{cases} g_\gamma(x, -x_{\text{esc}}) & ; -x_{\text{esc}} < x < x_{\text{inj}} \\ g_\gamma(x, -x_{\text{esc}}) - g_\gamma(x, x_{\text{inj}}) & ; x_{\text{inj}} < x < x_{\text{esc}}, \end{cases} \quad (21)$$

where $g_\gamma(x, y) = (A/\gamma)f_\gamma(x_{\text{esc}}, x_{\text{inj}})f_\gamma(x, y)/f_\gamma(x_{\text{esc}}, y)$ and $f_\gamma(x, y) = 1 - \exp[-\frac{2\pi\gamma}{3a}(x^3 - y^3)]$. The solution is extremely sensitive to $\gamma x_{\text{esc}}/a$. For $\gamma \gg \gamma_{\text{crit}} \equiv 3a/(2\pi x_{\text{esc}}^3)$, a broad wing of amplitude A/γ redward of x_{inj} will result as photons are redshifted across the line centre. The distribution will cut off sharply at $x > x_{\text{inj}}$. For an absorber with $T_a = 100$ K ($a \simeq 0.00472$), and $x_{\text{esc}} = 200$, $\gamma_{\text{crit}} \simeq 2.8 \times 10^{-10}$. The solutions for $\gamma = 1.7 \times 10^{-7}$ with $(x_{\text{inj}}, x_{\text{esc}}) = (100, 200)$ and (25, 50) are shown in Fig. 3. A substantial decrease in $n(x)$ is produced compared with the $\gamma = 0$ case. A significant enhancement of the Ly α collision rate over the free-streaming limit, however, may still result, with $P_{i,\text{ss}}/P_i(0) \simeq a/(\pi\gamma x_1)$. For $\gamma < 0$, a broad wing blueward of x_{inj} will form as photons are blueshifted in the contracting gas. A complex profile, however, may form redward of x_{inj} as redward diffusion is resisted by blueshifting. Such a case is illustrated in Fig. 3 for $\gamma = -1.7 \times 10^{-7}$ with $x_{\text{inj}} = 25$ and $x_{\text{esc}} = 50$.

3.2 Steady-state solution with atomic recoil

The above solutions neglect the effect of atomic recoil on the evolution of the radiation field. Since the photon momentum ($\sim h_P\nu_0/c$) is small compared with the momentum of the hydrogen atoms ($\sim m_H b$), the effect should be small, but may result in a significant heating rate when the gas is still cold (Madau et al. 1997; Chen & Miralda-Escudé 2004; Meiksin 2006). Allowing for atomic recoils modifies Eq. (14) to

$$\frac{1}{2} \frac{d}{dx} \left[2\epsilon\phi_V(a, x)n(x) + D(a, x)\frac{dn(x)}{dx} \right] = -\tilde{S}(x), \quad (22)$$

where the recoil parameter $\epsilon = h_P\nu_0/(2k_B T m_H c^2)^{1/2} \simeq 0.025 T^{-1/2}$ is the ratio of the momentum of a resonance line photon to the momentum of an atom moving at the thermal velocity (Rybicki & dell’Antonio 1994; Meiksin 2006). Eq. (22) expresses only the first order correction in ϵ . The solution for a source $\tilde{S}(x) = A\delta_D(x - x_{\text{inj}})$ is

$$n(x) = \begin{cases} g_\epsilon(x, -x_{\text{esc}}) & ; -x_{\text{esc}} < x < x_{\text{inj}} \\ g_\epsilon(x, -x_{\text{esc}}) - g_\epsilon(x, x_{\text{inj}}) & ; x_{\text{inj}} < x < x_{\text{esc}}, \end{cases} \quad (23)$$

where $g_\epsilon(x, y) = A[\pi/(2\epsilon^3 a)] \exp(-2\epsilon x)[f_\epsilon(x) - f_\epsilon(y)][f_\epsilon(x_{\text{esc}}) - f_\epsilon(x_{\text{inj}})]/[f_\epsilon(x_{\text{esc}}) - f_\epsilon(y)]$, and $f_\epsilon(x) = (1 - 2\epsilon x + 2\epsilon^2 x^2) \exp(2\epsilon x)$. Here, the wing approximation for $\phi_V(a, x)$ is adopted, but we find a nearly identical result allowing for the Doppler core as well.

The solution for $x_{\text{inj}} = 100$, $x_{\text{esc}} = 200$ and $\epsilon = 0.0079$, corresponding to $T \simeq 10$ K, is shown in Fig. 4. The result differs substantially from the solution for $\epsilon = 0$. In fact the solution cannot be trusted, as it is not consistent with the original order of Eq. (22) in ϵ . This may be demonstrated by adding to $2\epsilon\phi_V(a, x)n(x)$ the quantity $\epsilon^2 y(x)$, where $y(x)$ is an unspecified function representing the higher order corrections to the recoil, and which in general may also depend on $n(x)$. Inserting the series solution $n(x) = \sum_{i=0}^{\infty} \epsilon^i n_i(x)$ into Eq. (22) and equating equal orders in ϵ recovers $n_0(x)$ as given by Eq. (18), and produces the first order equation $(d/dx)[2\phi_V(a, x)n_0(x) + D(a, x)dn_1(x)/dx] = 0$ and the second order equation $(d/dx)[2\phi_V(a, x)n_1(x) + y(x) + D(a, x)dn_2(x)/dx] = 0$. The first order equation gives $n_1(x)$ in terms of $n_0(x)$, subject to the boundary conditions $n_1(-x_{\text{esc}}) = n_1(x_{\text{esc}}) = 0$. The second order equation shows that $n_2(x)$ depends on the unspecified (neglected) function $y(x)$. Thus the solution is valid only up to $n(x) = n_0(x) + \epsilon n_1(x)$. If higher order terms are substantial, then the solution for $n(x)$ cannot be trusted. The solution $n_0(x) + \epsilon n_1(x)$ for the example above differs substantially from the solution $n(x)$, and indeed is not even positive-definite, as shown in Fig. 4, demonstrating the solution is not valid.

Including higher order terms involves a non-trivial expression for the redistribution function including non-linear terms in the recoil parameter (Basko 1981; Meiksin 2006). (Additional higher order terms in the recoil parameter arise from relativistic corrections, but these are reduced by factors of b/c .) For temperatures $T > 100$ K, the discrepancy between the full solution and the first order solution is smaller than about 20% for $x_{\text{inj}} = 100$ and $x_{\text{esc}} = 200$. In general, the correction will be small for $\epsilon x_{\text{esc}} \ll 1$, and we caution that if this is violated the steady-state solutions may not be valid. We do not pursue further consequences of the atomic recoil solution here.

3.3 The role of bulk peculiar velocities

Absorption systems with a bulk peculiar velocity component away from the source will see the photons redshifted into line centre. In this case, the scattering rate of resonance line photons may be appreciable even in an optically thin absorber ($N_{19} < N_{19, \text{crit}}$), so that the build-up of a strong radiation field at line centre through photon capture and




Figure 5. Surface of constant Ly α scattering rate P_l produced by a continuum source to the left within a collapsing halo of neutral hydrogen. The inner surface terminates at the accretion shock. The circle represents the turnaround radius r_{ta} of the halo, with the solid line representing the hemisphere of gas with peculiar velocity redshifted relative to the source and the dashed line the hemisphere with blueshifted peculiar velocity. The surface shown corresponds to a maximum radius of $0.59r_{\text{ta}}$. The figure is tilted by 5° to reveal the 3D structure of the surface.

diffusion is not necessary for significant WFE decoupling of the spin state from the CMB. The peculiar velocity required to bring the absorber into the peak of the source function $S(\nu)$ in Eq. (16) is given by $v_{\text{pec}} \simeq \frac{1}{2}x_1b_a$, where x_1 is given by Eq. (4). For pure thermal broadening (assuming negligible internal velocity structure within the absorber), this corresponds to

$$v_{\text{pec}} \simeq 14h^{-1}(1+z)^{3/2} \left(\frac{T_a}{T_{\text{IGM}}} \right)^{1/2} \text{ km s}^{-1}. \quad (24)$$

This is comparable to the typical peculiar velocities of non-linear cosmological structures at the epochs of interest. As a result, the WFE will produce a 21cm signature that maps the peculiar velocity field of the structures within the IGM.

As an illustration, we consider the signal produced by a collapsing halo, such as around a forming galaxy or galaxy cluster, illuminated by an external radiation source, as depicted in Fig. 5.

The collision rate from a source of specific luminosity L_ν on an absorber with Doppler parameter b_a at luminosity distance r_L moving away from the source with peculiar velocity v is

$$\begin{aligned} P_l &= \sigma \frac{L_\nu}{h_P \nu} \frac{1}{4\pi r_L^2} \int_0^\infty d\nu \varphi_V \left[a, \nu \left(1 - \frac{v}{c} \right) \right] \exp(-\tau_\nu) \\ &\simeq \sigma \frac{L_\nu}{h_P \nu} \frac{1}{4\pi r_L^2} \exp \left[-\frac{b_a x_1}{v_0 \cos(\theta)} \right], \end{aligned} \quad (25)$$

using Eq. (3), approximating $\phi_V(a, x) \simeq \pi^{-1/2} \exp(-x^2)$, noting that the resulting exponential in the integrand peaks sharply near $x = v/b_a$ assuming $v_0/b_a \simeq x_1$ or larger, Taylor expanding the argument of the exponential to second order in $x - v/b_a$, and expressing the peculiar velocity as $v =$

$v_0 \cos(\theta)$, the projected accretion velocity along the line of sight to the source.

In general, v_0 will vary with distance from the centre of the collapsing halo. As a definite case, we approximate the peculiar velocity of the accreting gas using the self-similar solution of Bertschinger (1985) for adiabatic accretion of a $\gamma = 5/3$ gas of negligible density onto a dark matter halo with a comoving centre in an Einstein-de Sitter cosmology. The velocity field in the region between the accretion shock and the turn-around radius at cosmological time t is well-approximated by

$$v_0(r) \simeq \frac{r_{\text{ta}}}{t} \frac{V_{\text{min}}}{\log_{10} \lambda_s} \log_{10} \lambda, \quad (26)$$

where r_{ta} is the turn-around radius, $\lambda = r/r_{\text{ta}}$, $\lambda_s = r_s/r_{\text{ta}} \simeq 0.3472$, where r_s is the radius of the accretion shock, and $V_{\text{min}} \simeq -1.433$ characterises the inflow velocity of the gas just before passing through the accretion shock.

Contours of constant values of P_l correspond to values of constant projected accretion velocity. If at a radius r towards the source ($\theta = 0$), the accretion velocity is a fraction f of its maximum infall velocity, then contours of fixed P_l will correspond to the surfaces

$$\lambda = \lambda_s^{f \sec(\theta)}, \quad (27)$$

extending over the angular range $0 \leq \theta \leq \arccos(f)$. The surface corresponding to $f = 0.5$ is shown in Fig. 5, with a maximum angle $\theta_{\text{max}} = 60^\circ$ and maximum radius $\lambda_0 = \lambda_s^f = 0.59$. The loci of constant P_l correspond to arcing sheets. The exponential sensitivity of P_l to v for $v_0/b_a \simeq x_1$ (see Eq. [25]) ensures a surface of narrow width will dominate the 21cm signal produced. Such a surface could be mistaken for the region of neutral hydrogen just outside an H II region surrounding a source of ionizing radiation. A varying gas temperature, and therefore varying b_a , in the accreting region will further complicate the signal. If $v_0/b_a \gg x_1$, then the entire solid hemisphere will be illuminated by the source (except for a thin circular wafer at $\theta \lesssim 90^\circ$).

A similar effect may apply to an expanding void. Since voids expand faster than the Hubble expansion, the peculiar velocity on the far side of a void from a distant source could redshift the received radiation into line centre. A shell of direct scattering may then result even without including the reddening effects of radiative transfer within the void.

4 THE EVOLUTION OF THE WFE IN DISCRETE OBJECTS

It was assumed for the solutions in § 3 that the photons are able to achieve a steady state as they scatter within the absorbing system. In this section, we compute the timescales required to establish a steady state by solving the time-dependent diffusion equation

$$\frac{\partial n(x, \tau)}{\partial \tau} = \frac{1}{2} \frac{\partial}{\partial x} \left[D(a, x) \frac{\partial n(x, \tau)}{\partial x} \right] + \tilde{S}(x), \quad (28)$$

where $\tau = t/t_s$, and $t_s = \Delta\nu_D/(n_l \sigma c) \simeq 3.2n_l^{-1}T_a^{1/2}$ s is the scattering time of the resonance line photons at line centre.

We solve Eq. (28) using a Crank-Nicholson scheme. We tested the scheme against the time-dependent solutions for a




Figure 6. (a) Convergence of the time-dependent solution to the diffusion equation for a source $\tilde{S}(x) = \delta_D(x - x_{\text{inj}})$ with $x_{\text{inj}} = 100$, $x_{\text{esc}} = 200$ and $T_a = 100$ K. Shown are the steady-state photon frequency distribution, Eq. (18) (solid line), and the photon frequency distributions at the times their values at $x = 0$ take on 50% (dashed line), 90% (dotted line) and 97% (dot-dashed line) of the steady-state value. (b) The evolution of the ratio of the photon frequency distribution at line centre ($x = 0$) to the steady-state value, as a function of time in units of the scattering time at line centre.




Figure 7. The convergence time in units of the scattering time at line centre for the photon frequency distribution with source $\tilde{S}(x) = \delta_D(x - x_{\text{inj}})$ having $x_{\text{inj}} = x_{\text{esc}}/2$ to reach 50% (dashed line) and 90% (solid line) of the steady-state value at $x = 0$. A temperature of $T_a = 100$ K is assumed.

constant δ -function source at line centre in an infinite homogeneous medium and a flat continuum source in the diffusion approximation (Rybicki & dell’Antonio 1994). The results applied to the source $\tilde{S}(x) = \delta_D(x - x_{\text{inj}})$ with $x_{\text{inj}} = 100$ and boundary condition $n(x) = 0$ for $|x| > x_{\text{esc}} = 200$ are shown in the upper panel of Fig. 6. The photon frequency distribution takes on the approximate shape of the final steady-state solution Eq. (18) as it converges towards it. The evolution of the photon frequency distribution at line centre ($x = 0$) is shown in the lower panel of Fig. 6. For sources that have been active only a small fraction of the time required to achieve full convergence, the scattering rate may be negligibly small.

Dimensional analysis of Eq. (28) suggests the time to establish a steady state for photons injected at $x = x_{\text{inj}}$ is $\tau_{ss} \simeq 2x_{\text{inj}}^2/D(a, x_{\text{inj}}) \simeq (2\pi/a)x_{\text{inj}}^4 = (\pi/8a)x_{\text{esc}}^4 \simeq 8.3T_a^{1/2}x_{\text{esc}}^4$ for $x_{\text{inj}} = x_{\text{esc}}/2$. The time for the photon distribution to converge to the steady-state value at $x = 0$ found from the numerical integration of Eq. (28) is shown in Fig. 7 for convergence to 50% and 90% of the steady-state value, for a source $\tilde{S}(x) = \delta_D(x - x_{\text{inj}})$ with $x_{\text{inj}} = x_{\text{esc}}/2$ and $T_a = 100$ K. The convergence times scale like $\tau_{ss,50} \simeq 210x_{\text{esc}}^4$ (50%) and $\tau_{ss,90} \simeq 750x_{\text{esc}}^4$ (90%), in good agreement with the expected behaviour.

5 DISCUSSION AND CONCLUSIONS

The role played by the Wouthuysen-Field effect in decoupling the spin temperature of the neutral hydrogen from the CMB temperature depends most crucially on the scattering rate of Ly α photons. The spin temperature T_S is generally a weighted mean of the colour temperature T_α , kinetic temperature T_K of the gas, and brightness temperature T_R of any incident radiation (such as the CMB) at the 21cm frequency,

$$T_S = \frac{T_R + y_\alpha T_\alpha + y_c T_K}{1 + y_\alpha + y_c}, \quad (29)$$

where

$$y_\alpha \equiv \frac{P_{10}}{A_{10}} \frac{T_*}{T_\alpha} \quad \text{and} \quad y_c \equiv \frac{C_{10}}{A_{10}} \frac{T_*}{T_K} \quad (30)$$

are the Ly α and collisional pumping efficiencies, respectively (Field 1958; Madau et al. 1997), $A_{10} = 2.85 \times 10^{-15} \text{ s}^{-1}$ is the spontaneous decay rate of the 21cm transition, and $T_* \equiv h\nu_{10}/k_B$, where ν_{10} is the frequency of the 21cm transition. Here $P_{10} = 4P_l/27$ is the indirect de-excitation rate of the triplet hyperfine state induced by Ly α photon scattering. The colour temperature of the radiation field is the harmonic mean temperature $T_\alpha = 1/\langle T_u^{-1}(\nu) \rangle$, where $T_u(\nu) = -(h\nu/k_B)(d \ln u_\nu/d\nu)^{-1}$, weighted by $u_\nu \varphi_V(a, \nu)$ for an energy density u_ν of resonance line photons (Meiksin 2006). The coefficient C_{10} is the de-excitation rate of the triplet hyperfine state induced by atomic collisions.

For a Ly α scattering rate P_l exceeding the critical thermalization rate

$$P_{\text{th}} = \frac{27A_{10}T_{\text{CMB}}}{4T_*} \simeq 6.8 \times 10^{-12} \text{ s}^{-1} \left(\frac{1+z}{9} \right), \quad (31)$$

where $T_{\text{CMB}} \simeq 2.73(1+z)$ (Mather & et al. 1994) is the CMB temperature, the spin temperature will be driven to

the colour temperature in the absence of strong collisional de-excitation (Madau et al. 1997). The radiation field will typically rapidly thermalize with the neutral hydrogen, resulting in $T_\alpha \rightarrow T_K$ (Field 1959b; Meiksin 2006). The neutral hydrogen will then appear in either absorption or emission against the CMB, depending on whether T_K is below or above T_{CMB} , respectively.

Before the Epoch of Reionization, when the first radiation sources turn on at the End of the Dark Ages, regions around the sources may become visible through their 21cm signature. Because the photons produced by a source, such as a QSO or star-forming galaxy, near the $\text{Ly}\alpha$ resonance will be scattered out of the line of sight over short distances, the scattering rate of $\text{Ly}\alpha$ photons will be much smaller than P_{th} except extremely near the source. Only photons emitted sufficiently blueward of the $\text{Ly}\alpha$ resonance frequency will survive to any distant structures, where they will be received well in the blue Lorentz wing. If a structure has a sufficient optical depth in the wing to the photons received, it will trap them and rescatterings within the structure will allow the photons to diffuse in frequency across the resonance line. The radiation field may be built up in this way until it reaches a steady state in which the rate of incoming photons is balanced by the rate at which photons diffuse sufficiently far from the resonance line frequency to escape in the wings where the structure is optically thin. The resulting photon scattering rate in the diffusion approximation may greatly exceed the estimate $P_l(0)$ in the free-streaming limit, for which it is assumed no resonance line photons emitted by the source have been scattered out of the line of sight before reaching the absorber.

We identify several factors, however, that may reduce the energy density of the resonance line photons from the steady-state diffusion approximation estimate for static structures:

(1) Internal expansion within the structure, characterised by the dimensionless expansion parameter γ , will redshift the photons into a red wing, where they will escape, on a timescale much faster than the diffusion time, resulting in a much reduced energy density. In the diffusion approximation, the scattering rate will be reduced compared with the $\gamma = 0$ case by the factor $(3a/\pi\gamma)/(x_{\text{esc}}^3 - x_{\text{inj}}^3)$ for $\gamma \gg \gamma_{\text{crit}} = 3a/(2\pi x_{\text{esc}}^3)$, where x_{esc} is the dimensionless escape frequency at which photons escape the structure, and x_{inj} is the frequency at which photons are predominantly received from the source. Contraction within the structure may altogether prevent photons from diffusing across the resonance line frequency.

(2) The scattering rate may be reduced by atomic recoil. Estimating the magnitude of this effect, however, generally requires including non-trivial higher order terms in the recoil parameter ϵ than first order in the steady-state equation. We caution against introducing atomic recoils into Monte Carlo calculations and other steady-state solutions, as the results obtained may not be physically self-consistent if the solutions differ substantially from those obtained for $\epsilon = 0$.

(3) The time scale to establish a steady state will typically exceed the duration of high star formation rates in galaxies or the lifetime of QSOs. As a consequence, the collision rate will be reduced by factors of tens to thousands compared with the steady-state value.

As an illustration, consider the region around a bright QSO with specific luminosity at the Lyman edge of $10^{31} \text{ erg s}^{-1} \text{ Hz}^{-1}$ pre-heated by soft x-rays ahead of the ionization front to a mean temperature of $T_{\text{IGM}} \simeq 1800 \text{ K}$ at a distance of 10 Mpc (Madau et al. 1997). According to Eq. (4), $x_1 \simeq 200$ at $z = 8$ (assuming $h = 0.7$). A structure 250 kpc across with a moderate overdensity of 3 and internal temperature of $T_a \simeq 120 \text{ K}$ will have a neutral hydrogen column density of $N_{\text{HI}} \simeq 3 \times 10^{20} \text{ cm}^{-2}$ and escape frequency $x_{\text{esc}} \simeq 200$, from Eq. (12). For a power law spectrum $\nu^{-1.5}$, Eq. (10) gives for the $\text{Ly}\alpha$ collision rate in the free-streaming limit $P_l(0) \simeq 8.7 \times 10^{-13} \text{ s}^{-1}$, about an order of magnitude smaller than the thermalization rate P_{th} . The steady-state value from Eq. (19) then gives $P_{l,\text{ss}} \simeq 10^{-8} \text{ s}^{-1}$, well in excess of P_{th} . The time to establish a steady state, however, is long. The scattering time of $\text{Ly}\alpha$ photons in the structure is about $t_s \simeq 8.5 \times 10^4 \text{ s}$. According to Fig. 7, the time for the radiation field to achieve a steady state is then about 10^{17} s , or 3 Gyr, greatly exceeding the expected lifetime of the QSO. If the QSO had illuminated the structure for as long as 0.1 Gyr, according to Fig. 6 the steady-state value must be reduced by a factor of 0.05. The resulting scattering rate will still exceed P_{th} . If the system has a peculiar velocity away from the QSO, the rate could be quite different. Eq. (24) shows that a peculiar velocity of 140 km s^{-1} , a plausible value, would shift the peak of the radiation from the source received by the structure to the resonance line frequency. The radiation field would then rapidly reach its steady-state value, and $P_{l,\text{ss}}$ would greatly exceed P_{th} .

A much more overdense structure at the same distance from the QSO would be able to achieve the steady-state scattering rate. For example, a virialized structure 35 kpc across with an overdensity of 200 and temperature $T_a \simeq 1000 \text{ K}$ would have a neutral hydrogen column density of $3 \times 10^{21} \text{ cm}^{-2}$ and $x_{\text{esc}} \simeq 220$. The time for the radiation field to reach a steady state would then be about 0.2 Gyr, a plausible lifetime for the QSO.

By contrast, it may be difficult to achieve a steady state in a void. For an IGM temperature $T_{\text{IGM}} \simeq 100 \text{ K}$, $x_1 = 825$ at $z = 8$. A region 3 Mpc across with a temperature of $T_a = 10 \text{ K}$, and underdense by a factor of 3 will have a neutral hydrogen column density of $4 \times 10^{20} \text{ cm}^{-2}$, $x_{\text{esc}} \simeq 800$ and a scattering time at line centre of $t_s \simeq 2.2 \times 10^5 \text{ s}$. The time to reach a steady state is then 2100 Gyr. Even if the void is illuminated for as long as the age of the Universe, the radiation field will be negligibly small compared with its steady-state value. Since the void will expand, however, the incident radiation will be redshifted across the resonance line. Rybicki & dell’Antonio (1994) estimate a characteristic time for redshifting to dominate diffusion in an infinite homogeneous medium to be $\tau_\gamma \simeq (a/\gamma^4)^{1/3}$. Voids expand somewhat faster than the mean Hubble expansion. Adopting the Hubble constant at $z = 8$ for $h = 0.7$, $\gamma \simeq 5 \times 10^{-6}$. A steady state should be established for $t \gg \tau_\gamma t_s \simeq 2 \times 10^4 \text{ yr}$, and the steady-state scattering rate will then match the free-streaming value, with $P_{l,\text{ss}}/P_l(0) \simeq a/(\pi\gamma x_1) \simeq 1$, which may be adequate for producing a 21cm signature in the presence of sufficient sources.

The examples above show that the effect of peculiar motions may lead to an enhancement in the $\text{Ly}\alpha$ scattering rate over the free-streaming limit and produce the required decoupling of T_S from T_{CMB} . Even if the scattering rate is

inadequate for inducing decoupling throughout most of the structure, it may be adequate along narrow loci within which the velocity field either permits the resonance line photons to accumulate near the resonance line frequency or shifts the resonance line frequency of the absorber to matching the incident radiation field from the source as filtered through the IGM. We presented an example of an accreting halo which produces a 21cm signature along a curved surface for which the impinging photons are received at the resonance line frequency due to the infall velocity of the accreting gas. Such a curved structure could be mistaken for a rim of neutral gas outside an H II region in radio maps of the IGM. Similar curved surfaces may form on the far side of voids from a distant source.

As a consequence of the above effects, the 21cm signature of the IGM will not simply trace the density structure of the IGM, but will trace the peculiar velocity structure as well. We have not considered multiple sources. In the presence of a large number of nearby sources, a greater fraction of the gas will be able to match the incident radiation field through their peculiar motions. In addition, the escaping radiation from the absorbers will contribute to the ambient radiation field. In due course, a metagalactic diffuse radiation field may grow sufficiently strong to ensure the WFE is capable of decoupling the spin temperature from the CMB temperature throughout the IGM, and so ensure the production of a 21cm signature everywhere. The effects described here suggest that in the early stages, before any such metagalactic field has been established (and it has yet to be demonstrated that such a field will be established), the 21cm signatures produced at the End of the Dark Ages, and possibly extending into the Epoch of Reionization, may only be interpreted with a thorough understanding of the evolution of the energy density of resonance line photons within a clumpy medium, including the effects of peculiar motions.

APPENDIX A: OPTICAL DEPTH THROUGH DOPPLER CORE

Radiation from a source at z_S received at z with frequency ν in the range $\nu_0[(1+z)/(1+z_S)] < \nu < \nu_0$, where ν_0 is the resonance line frequency, will have passed through the resonance line frequency en route to the absorber. To estimate the optical depth of the IGM at these frequencies, it is convenient to approximate the Voigt line profile as

$$\begin{aligned}\phi_V(a, x) &= \pi^{-1/2} \exp(-x^2); & -x_m < x < x_m \\ &= \frac{a}{\pi x^2}; & |x| > x_m\end{aligned}\quad (A1)$$

where x_m is the value of $x = (\nu - \nu_0)/\Delta\nu_D$ at which the two approximations match. For $T = 10$ K, 100 K and 1000 K, the values of x_m are 2.58, 2.83 and 3.05, respectively.

A photon of frequency ν at z will have $x = \pm x_m$ at the redshift $z_m^{(\pm)}$, given by

$$\frac{\nu}{\nu_0} \frac{1+z_m^{(\pm)}}{1+z} = 1 \pm x_m \frac{b}{c} \quad (A2)$$

where $b = (2k_B T_{\text{IGM}}/m_H)^{1/2}$ is the Doppler parameter of the IGM at temperature T_{IGM} , assumed constant. (In principle, a turbulent velocity component may be added in

quadrature.) The optical depth in Eq. (1) may then be expressed as a sum of three contributions, corresponding to the ranges $x < -x_m$, $-x_m < x < x_m$ and $x > x_m$,

$$\begin{aligned}\tau_\nu &= \frac{\sigma a}{\pi \Delta\nu_D} \int_z^{z_m^{(-)}} dz' \frac{dl_p}{dz'} n_l(z') \frac{(\Delta\nu_D)^2}{\left(\nu \frac{1+z'}{1+z} - \nu_0\right)^2} \\ &+ \frac{\sigma}{\pi^{1/2} \Delta\nu_D} \int_{z_m^{(-)}}^{z_m^{(+)}} dz' \frac{dl_p}{dz'} n_l(z') \exp \left[- \left(\frac{\nu \frac{1+z'}{1+z} - \nu_0}{\Delta\nu_D} \right)^2 \right] \\ &+ \frac{\sigma a}{\pi \Delta\nu_D} \int_{z_m^{(+)}}^{z_S} dz' \frac{dl_p}{dz'} n_l(z') \frac{(\Delta\nu_D)^2}{\left(\nu \frac{1+z'}{1+z} - \nu_0\right)^2}.\end{aligned}\quad (A3)$$

The first and third terms are similar to Eq. (2). The dominant term is the second, representing scattering through the line core. Setting $u = [(\nu(1+z')/(1+z) - \nu_0)/\Delta\nu_D]^2$, and assuming $b/c \ll 1$ within the integrand, gives for this contribution

$$\begin{aligned}\tau_\nu^{\text{core}} &\simeq \frac{1}{\pi^{1/2}} \frac{\sigma c n(0)}{H_0 \Omega_m^{1/2}} (1+z)^{3/2} \nu_0^{1/2} \nu^{-3/2} \gamma\left(\frac{1}{2}, x_m^2\right) \\ &\simeq 8000 h^{-1} (1+z)^{3/2} y^{-3/2} \gamma\left(\frac{1}{2}, x_m^2\right),\end{aligned}\quad (A4)$$

where $y = \nu/\nu_0$, $\gamma(t, x) = \int_0^x du u^{t-1} \exp(-u)$ is an incomplete gamma function, and $\Omega_m = 0.3$ and $\Omega_b h^2 = 0.022$ are assumed. This will in general ensure that essentially no flux from the source is received in the frequency range $\nu_0(1+z)/(1+z_S) < \nu < \nu_0$. This is the Gunn-Peterson effect.

APPENDIX B: SLAB SOLUTION

The diffusion equation for the transfer of radiation through a static slab in the Eddington approximation is given by Harrington (1973) as

$$\begin{aligned}\frac{\partial^2 n_x(\tau, \sigma)}{\partial \tau^2} + \frac{\partial^2 n_x(\tau, \sigma)}{\partial \sigma^2} &= -3\phi_V^2(a, x) \frac{G(\tau)}{4\pi}, \\ &\simeq -6^{1/2} \delta_D(\sigma) \frac{G(\tau)}{4\pi},\end{aligned}\quad (B1)$$

for a spectrally flat source $G(\tau)$, in the limit $\phi_V(a, x) \simeq a/(\pi x^2)$. Here, σ is related to x through $dx/d\sigma = (3/2)^{1/2} \phi_V(a, x)$, and $\tau = \tau_\nu/\phi_V(a, x)$ is the mean optical depth vertically through the slab. The photon number density $n(x)$ is expressed as $n_x(\sigma)$. The last line in Eq. (B1) follows from approximating $3\phi_V^2(a, x)$ as a Dirac δ -function, $6^{1/2} \delta_D(\sigma)$ where the coefficient preserves the normalization. The boundary conditions assumed are

$$\left. \frac{\partial n_x(\tau, \sigma)}{\partial \tau} \right|_{\pm B} = \mp \frac{3}{2} \phi_V(a, x) n_x(B, \sigma) \quad (B2)$$

and

$$n_x(\tau, \sigma) \rightarrow 0 \quad \text{for } \sigma \rightarrow \pm\infty. \quad (B3)$$

For a uniformly distributed source of unit strength, Harrington (1973) takes $G = 1/(2B)$, where $-B < \tau < B$ describes the vertical extent of the slab. In terms of the total optical depth at line centre, $\tau_0 = 2B\phi_V(a, 0) = 2B/\pi^{1/2}$, the source becomes $G = \pi^{-1/2}/\tau_0$. The solution at the centre of the slab is

$$n_x(0, x) = \frac{6^{1/2}}{2\pi^3} S \left[\exp \left(-2^{1/2} \left(\frac{\pi}{3} \right)^{3/2} \frac{|x|^3}{a\tau_0} \right) \right], \quad (\text{B4})$$

where the function $S(z) = z - z^3/3^2 + z^5/5^2 - \dots \simeq z$ for $0 < z < 1$.

The diffusion approximation Eq. (14) transforms to

$$\frac{d^2 n_x(\sigma)}{d\sigma^2} = -6^{1/2} A \delta_D(\sigma), \quad (\text{B5})$$

for a source $\tilde{S}(x) = A\delta_D(x)$ and adopting $D(a, x) = \phi_V(a, x)$. Identifying $A = G/(4\pi) = 1/(4\pi^{3/2}\tau_0)$ reproduces Eq. (B1) with the τ -dependence suppressed. Expressing $x_{\text{esc}} = x_*(a\tau_0)^{1/3}$, the solution Eq. (18) becomes

$$n_x(x) = \frac{x_*^3}{12\pi^{1/2}} \left(1 - \frac{|x|^3}{x_*^3 a\tau_0} \right) \quad (|x| \leq x_{\text{esc}}), \quad (\text{B6})$$

for $x_{\text{inj}} = 0$. Noting $S(1) \simeq 0.92$, equating the values of Eqs. (B4) and (B6) at $x = 0$ gives $x_* \simeq 0.92$. The two profiles are very similar, with Eq. (B4) forming an exponential tail with a cutoff at $x \gtrsim 0.87(a\tau_0)^{1/3} \simeq 0.95x_{\text{esc}}$.

REFERENCES

- Adams T. F., 1972, ApJ, 174, 439
 Basko M. M., 1981, Astrophysics, 17, 69
 Becker R. H., et al. 2001, AJ, 122, 2850
 Bertschinger E., 1985, ApJS, 58, 39
 Chen X., Miralda-Escudé J., 2004, ApJ, 602, 1, arXiv:astro-ph/0303395
 Chuzhoy L., Zheng Z., 2007, ApJ, 670, 912, arXiv:0706.0895
 Fan X., Carilli C. L., Keating B., 2006, Ann. Rev. Astron. Astrophys., 44, 415, arXiv:astro-ph/0602375
 Field G. B., 1958, Proc. I.R.E., 46, 240
 Field G. B., 1959a, ApJ, 129, 536
 Field G. B., 1959b, ApJ, 129, 551
 Furlanetto S. R., Oh S. P., Briggs F. H., 2006, Phys. Rep., 433, 181, arXiv:astro-ph/0608032
 Gnedin N. Y., Shaver P. A., 2004, ApJ, 608, 611, arXiv:astro-ph/0312005
 Gunn J. E., Peterson B. A., 1965, ApJ, 142, 1633
 Harrington J. P., 1973, MNRAS, 162, 43
 Hogan C. J., Rees M. J., 1979, MNRAS, 188, 791
 Loeb A., Barkana R., 2001, Ann. Rev. Astron. Astrophys., 39, 19, arXiv:astro-ph/0010467
 Loeb A., Rybicki G. B., 1999, ApJ, 524, 527, arXiv:astro-ph/9902180
 Madau P., Meiksin A., Rees M. J., 1997, ApJ, 475, 429
 Mather J. C., et al. 1994, ApJ, 420, 439
 Meiksin A., 2006, MNRAS, 370, 2025
 Mihalas D., 1978, Stellar atmospheres /2nd edition/. San Francisco, W. H. Freeman and Co., 1978. 650 p.
 Neufeld D. A., 1990, ApJ, 350, 216
 O'Meara J. M., Burles S., Prochaska J. X., Prochter G. E., Bernstein R. A., Burgess K. M., 2006, ApJ, 649, L61, arXiv:astro-ph/0608302
 Osterbrock D. E., 1962, ApJ, 135, 195
 Pierleoni M., Maselli A., Ciardi B., 2007, ArXiv e-prints, 712, 0712.1159
 Rybicki G. B., dell'Antonio I. P., 1994, ApJ, 427, 603, arXiv:astro-ph/9312006
 Scott D., Rees M. J., 1990, MNRAS, 247, 510
 Semelin B., Combes F., Baek S., 2007, A&Ap, 474, 365, arXiv:0707.2483
 Spergel D. N., et al. 2007, ApJS, 170, 377, arXiv:astro-ph/0603449
 Tozzi P., Madau P., Meiksin A., Rees M. J., 2000, ApJ, 528, 597, arXiv:astro-ph/9903139
 Wouthuysen S. A., 1952, AJ, 57, 31
 Zhang Y., Meiksin A., Anninos P., Norman M. L., 1998, ApJ, 495, 63